

Springer Proceedings in Complexity

Samsul Ariffin Abdul Karim ·
Mohd Fadhlullah Abd Shukur ·
Chong Fai Kait · Hassan Soleimani ·
Hamzah Sakidin *Editors*

Proceedings of the 6th International Conference on Fundamental and Applied Sciences

ICFAS 2020

 Springer



Conference proceedings | © 2021

Proceedings of the 6th International Conference on Fundamental and Applied Sciences

ICFAS 2020

[Home](#) > Conference proceedings

Editors: [Samsul Ariffin Abdul Karim](#), [Mohd Fadhlullah Abd Shukur](#), [Chong Fai Kait](#), [Hassan Soleimani](#), [Hamzah Sakidin](#)

Presents peer-reviewed articles from ICFAS 2020

Brings together leading experts, scientific communities in the field of applied sciences and mathematics

Includes topics such as green materials, molecular modeling, nanodevices

Part of the book series: [Springer Proceedings in Complexity](#) (SPCOM)

17k Accesses | **19** Citations | **1** Altmetric

Sections

[Table of contents](#)[About this book](#)[Keywords](#)[Editors and Affiliations](#)[About the editors](#)[Bibliographic Information](#)

This is a preview of subscription content, [access via your institution.](#)

Table of contents (67 papers)

Search within book

[← Previous](#)

Page

1

of 4

[Next →](#)**Front Matter**[PDF ↓](#)

Pages i-xxxii

Sustainable Chemistry

Front Matter[PDF ↓](#)

Pages 1-1

[Acoustic Performance of Mixing EFB and OPF Low-Density Fibreboards in Different Thickness](#)

Mageswaran Ravi Chandran, Ewe Lay Sheng,
Mohammad Nazhan Nasir, Yew Weng Kean, Zawawi
Ibrahim

Pages 3-13

[Modified Alginate Adsorbent Using Ionic Liquid for Manganese Removal from Aqueous Solution](#)

Noor A.'in A. Rahman, Gowri Selvaraj, Cecilia Devi
Wilfred
Pages 15-25

[Synthesis and Modification of Pour Point
Depressant \(PPD\) Based on Copolymers of
 \$\alpha\$ -Olefins and Maleic Anhydride for Waxy
Crude Oil](#)

Kozhabekov Serik Samsalykovich, Zhubanov Amin
Abdirasululy, Donenov Beisen Kainarbaevich,
Makhmetova Aliya Ruslanovna, Abayev Talgat Bakytuly
Pages 27-35

[Effects of NPK Fertilizers on the Growth,
Yield and Chemical Content of Tomato
\(*Lycopersicon esculentum* L. Mill\)](#)

Lyazzat Bekbayeva, Zuraini Zakaria, Tatyana Karpenyuk,
Alla Goncharova, El-Sayed Negim, Kozhanova Kaldanay
Pages 37-50

[Production of Greener Biodiesel from a
Low-Cost CaO Catalysts of Waste
Freshwater Shells](#)

Nazrizawati Ahmad Tajuddin, Nurul Natasha Shahrom,
Noraini Hamzah
Pages 51-60

[CO₂ Hydrogenation to Methanol: Effects of
Reaction Temperatures and Pellet Crushing
on the Catalytic Performance](#)

Nor Hafizah Berahim, Noor Asmawati Mohd Zabidi,
Nadia Syazana Abd Halim, Akbar Abu Seman, Nor
Hafizah Yasin
Pages 61-69

[Catalytic Conversion of CO₂ into Alcohols:
Comparison of Supports](#)

Nur Insyirah Zulkifli, Noor Asmawati Mohd Zabidi
Pages 71-84

[Raman Spectroscopy of TiO₂ Nanoparticles Synthesized by Hydrolysis of TiCl₄: Effect of Sulfate Ions Concentration](#)

Kar Mun Lee, Chong Fai Kait, Jun Wei Lim, Geok Bee Teh
Pages 85-95

[Growth of Lactobacillus casei and Propionibacterium jensenii in Different Glucose Concentration and Incubation Temperature](#)

S. M. Mohamed Esivan, R. Rashid, A. Jati, N. A. Zaharudin
Pages 97-105

[A Comparison of Two Methods for the Development of Low-Cost Carbonaceous Adsorbent from Rubber-Seed Shell \(RSS\)](#)

Syeda Saba Fatima, Azry Borhan, Muhammad Faheem
Pages 107-116

[Adsorptive Removal of Bisphenol A Using Zeolitic Imidazolate Framework \(ZIF-8\)](#)

Afzan Mahmad, Maizatul Shima Shaharun, Zakariyya Uba Zango, Teh Ubaidah Noh, Bahruddin Saad
Pages 117-129

[Inclusion Complexes of \$\beta\$ -Cyclodextrin/Pioglitazone and \$\beta\$ -Cyclodextrin-Ionic Liquid/Pioglitazone: Spectroscopic Methods Combined with Molecular Docking Studies](#)

Nurul Yani Rahim, Nur Najihah Md Zuki, Khairulazhar Jumbri, Sharifah Mohamad, Ninie Suhana Abdul Manan, Nadiyah Sidek et al.
Pages 131-145

[Mechanism of Alkaline Surfactant Polymer in Oil-Water Interface: Physicochemical of Fluid Rheology, Interfacial Tension and Molecular Dynamics Simulation](#)

Mohd Sofi Numin, Khairulazhar Jumbri, Anita Ramli,
Noorazlenawati Borhan
Pages 147-155

[Free Energy Calculation of CO₂ in Cholinium-Based Amino Acid Ionic Liquids](#)

Fitri Norizatit Salehin, Khairulazhar Jumbri, Anita Ramli,
Mohd Azlan Kassim, Mohd Faisal Taha
Pages 157-173

[Microwave-Assisted Solvothermal Liquefaction of Kenaf Stalks for Bio-oil Production](#)

Anita Ramli, Lai Weng Kin, Nur Akila Syakida Idayu
Khairul Anuar, Normawati M. Yunus, Nurul Jannah Abd
Rahman
Pages 175-184

[Interaction of TFSI-Imidazole and TFSI-Pyridinium ILs with MOFs from Molecular Docking Simulation](#)

Nor Ain Fathihah Abdullah, Khairulazhar Jumbri, Nurul
Yani Rahim
Pages 185-194

[Impact of Photoperiod on the Carbon Metabolic Pathways of Chlorella Vulgaris for Biomass Production and Nutrient Removal in Treating Nutrient-Rich Wastewater](#)

Nur Afiqah Mohamad Saman, Wai Hong Leong,
Hemamalini Rawindran, Mardawani Mohamad, **Muslim
Abdurrahman**, Jun Wei Lim
Pages 195-202

[Effect of Anodization Voltage on TiO₂ Nanotubes for Photodegradation of Formaldehyde](#)

Nurul Tasnim Sahrin, Chong Fai Kait, Lee Siew Ling,
Mohd Dzul Hakim Wirzal
Pages 203-212

[← Previous](#)

Page

1

of 4

[Next →](#)[Back to top ↑](#)

About this book

This book highlights latest advancement in Mathematics, Physics and Chemistry. With the theme of "***Innovative Science towards Sustainability and Industrial Revolution 4.0***", ICFAS 2020 brings together leading experts, scientific communities and industrialists working in the field of applied sciences and mathematics from all over the world to share the most recent developments and cutting-edge discoveries addressing sustainability and industrial revolution 4.0 in the field. The conference topics include green materials, molecular modelling, catalysis, nanodevices and nanosystems, smart materials applications, solar cells technology, computational mathematics, data analysis and visualization, and numerical analysis. The contents of this book are useful for researchers, students, and industrial practitioners in the areas of Mathematics, Physics and Chemistry as most of the topics are in line with IR 4.0.

[Back to top ↑](#)

Keywords

Sustainable chemistry**Nanodevices and Nanosystems****Smart materials Fuel cell technologies**

Numerical analysis **Molecular modeling**

Green materials **Bioremediation**

[Back to top](#) ↑

Editors and Affiliations

Fundamental and Applied Sciences, Universiti Teknologi PETRONAS, Seri Iskandar, Malaysia

Samsul Ariffin Abdul Karim, Mohd
Fadhlullah Abd Shukur

Universiti Teknologi PETRONAS, Seri Iskandar, Malaysia

Chong Fai Kait, Hassan Soleimani, Hamzah
Sakidin

[Back to top](#) ↑

About the editors

Samsul Ariffin Abdul Karim is a Senior Lecturer at Universiti Teknologi PETRONAS, Malaysia. He is a Professional Technologists registered with Malaysia Board of Technologists (MBOT). He has published more than 140 papers in Journals and Conferences including three Edited Conferences Volume and 40 book chapters. He was the recipient of Effective Education Delivery Award and Publication Award (Journal & Conference Paper), UTP Quality Day 2010, 2011 and 2012, respectively. He was Certified WOLFRAM Technology Associate, Mathematica Student Level. He also has published nine books with Springer Publishing including five books with Studies in Systems, Decision and Control (SSDC) series.

Muhammad Fadhlullah Abd. Shukur is a Senior Lecturer at Universiti Teknologi PETRONAS, Malaysia. His research interest is in the field of ionic conductor including the development and characterization of polymer electrolytes for application in various energy storage devices. He has received several research grants, acting as the head researcher or co-researcher for such grants. He has been invited as reviewer for various journals and has published 41 articles in ISI/SCOPUS indexed journals. Currently, his H-index is 21 with 1347 citations in Google Scholar Citations.

Chong Fai Kait is an associate professor at Universiti Teknologi PETRONAS, Malaysia. She graduated with B.Sc. (Hons) in Industrial Chemistry from Universiti Putra Malaysia in 1993. In 1998, she received her Ph.D. from the University of Dundee, Scotland, in the area of heterogeneous catalysis for naphtha reforming process. She has more than 20 years of academic and research experience and her main research interests are in photocatalysis and its integration with ionic liquids or other techniques for sustainable environmental applications.

Hassan Soleimani is an associate professor of Universiti Teknologi PETRONAS, Malaysia. He has obtained a Ph.D. in wave propagation from Universiti Putra Malaysia. He has published more than 120 research articles in peer reviewed International. He is committee member and organizer of International conference of Diffusion in Solid and Liquids (DSL) since 2011 in Europe. He has been visiting scientist at many universities including University of Cambridge, UK, Wright State University, USA, University of Patras, Greece and School of Engineering and Griffith University, Australia.

Hamzah Bin Sakidin is a Senior Lecturer at Universiti Teknologi PETRONAS, Malaysia since September 2013. His research interest is in the field of applied mathematics and mathematical modelling. He has received several research grants, citing as the head researcher or co-researcher for some research grants. He is also the author of several mathematics book for tertiary level of education. He has more than 20 years

of experience teaching in secondary and tertiary level of education.

[Back to top](#) ↑

Bibliographic Information

Book Title	Book Subtitle	Editors
Proceedings of the 6th International Conference on Fundamental and Applied Sciences	ICFAS 2020	Samsul Ariffin Abdul Karim, Mohd Fadhlullah Abd Shukur, Chong Fai Kait, Hassan Soleimani, Hamzah Sakidin

Series Title	DOI	Publisher
Springer Proceedings in Complexity	https://doi.org/10.1007/978-981-16-4513-6	Springer Singapore

eBook Packages	Copyright Information	Hardcover ISBN
Physics and Astronomy, Physics and Astronomy (R0)	Institute of Technology PETRONAS Sdn. Bhd. 2021	978-981-16-4512-9 Published: 06 January 2022

Softcover ISBN	eBook ISBN	Series ISSN
978-981-16-4515-0 Published: 07 January 2023	978-981-16-4513-6 Published: 05 January 2022	2213-8684

Series E-ISSN	Edition Number	Number of Pages
2213-8692	1	XXXI, 782

Number of Illustrations	Topics
85 b/w illustrations, 253 illustrations in colour	Complex Systems, Green Chemistry, Numerical Analysis

[Nanoscale](#)
[Devices](#)

[Back to top](#) ↑

Not logged in - 103.148.198.139

Not affiliated

SPRINGER NATURE

© 2023 Springer Nature Switzerland AG. Part of [Springer Nature](#).

Impact of Photoperiod on the Carbon Metabolic Pathways of *Chlorella Vulgaris* for Biomass Production and Nutrient Removal in Treating Nutrient-Rich Wastewater



Nur Afiqah Mohamad Saman, Wai Hong Leong, Hemamalini Rawindran, Mardawani Mohamad, Muslim Abdurrahman, and Jun Wei Lim

Abstract Cultivation of *Chlorella vulgaris* has been done in different photoperiod to study the effect of photoperiod cycles on the on the carbon metabolic pathways of the microalgae species in order to enhance its biomass and lipid productions. Autotrophic, heterotrophic and three different photoperiod cycles of mixotrophic growth mode were evaluated for its biomass growth, carbon and nutrient uptakes and lipid yield. The studied photoperiods for mixotrophic growth are 16:8, 12:12 and 8:16 h (light:dark hours). Heterotrophic condition produced much lower microalgal biomass and lipid yield compared to microalgae grown under autotrophic and mixotrophic condition. There is no significant difference in microalgal biomass yields observed under continuous illumination between 16:8 h, 12:12 h and 8:16 h photoperiods. All studied parameters showed a near complete removal of COD on the second day of cultivation. Higher nitrogen removal was observed at longer photoperiod conditions.

Keywords Microalgae · Photoperiod · Biomass · Carbon · Nitrogen

N. A. M. Saman

Department of Fundamental and Applied Sciences, Universiti Teknologi PETRONAS, 32610 Seri Iskandar, Perak Darul Ridzuan, Malaysia
e-mail: nur.afiqah_25213@utp.edu.my

W. H. Leong · H. Rawindran · J. W. Lim (✉)

Department of Fundamental and Applied Sciences, HICoE—Centre for Biofuel and Biochemical Research, Institute of Self-Sustainable Building, Universiti Teknologi PETRONAS, 32610 Seri Iskandar, Perak Darul Ridzuan, Malaysia
e-mail: junwei.lim@utp.edu.my

M. Mohamad

Faculty of Bioengineering and Technology, Universiti Malaysia Kelantan, Jeli Campus, 17600 Jeli, Kelantan, Malaysia
e-mail: mardawani.m@umk.edu.my

M. Abdurrahman

Department of Petroleum Engineering, Universitas Islam Riau, Jl. Kaharuddin Nasution No. 113, Pekanbaru 28284, Indonesia
e-mail: muslim@eng.uir.ac.id

1 Introduction

Microalgae is steadily gaining attraction as the preferred feedstock for sustainable biofuel productions termed as the third-generation biofuels. Moreover, microalgae have the ability in fixing atmospheric carbon dioxide for its growth and produce a variety of end products [1]. Production rate of biodiesel from microalgae is about 5000–20,000 gal/acre/year which is comparatively high than oil production from other feedstocks [2]. In addition, some microalgae strains are very versatile that they can be grown in extreme environments and conditions e.g. high nutrient wastewater loadings and saline water bodies [3–5].

While the merits of using microalgae as a potential biofuel feedstock are undeniably promising, the commercialization of microalgae-based biofuels is rather uncertain [6]. Thus, leading to extensive studies to further enhance the biomass and lipid productivities for eventual exploitations as microalgal biomass feedstock. These include manipulation of cultivation parameters and cultivation modes. The main obstacle that impedes the commercialization of microalgal biodiesel production is the low lipid productivities due to the low cell biomass and lipid production. Owing to the low biomass yield, the harvesting process become difficult which can subsequently lead to high unit cost of downstream process. Various studies into optimizing or modifying the cultivation conditions have been investigated to induce both the microalgal biomass and lipid productivities [7, 8]. As microalgae are known to be able to utilize various carbon metabolic pathways, namely, autotrophic, heterotrophic and mixotrophic, for its growth, manipulating its growth metabolic pathways perhaps could provide an insight into inducing the cellular biomass and lipid productions [9].

Other than that, photoperiod is one of the important physicochemical factors affecting the growth of microalgae as it regulates the cell division in microalgae reproduction. The influence of this photoperiod on biomass and lipid productivity of microalgae has not been clearly established. To further improve the biomass yields of microalgae, it is crucial to optimize the photoperiod to achieve high microalgal biomass growth productivities. Hence, this study is aimed to investigate the carbon metabolic pathways through varying the light and dark cycles i.e., photoperiod for optimizing the microalgal biomass production along with the nutrient removal from nutrient-rich wastewater. Thus, able to provide an insight on the dynamics of microalgal biomass growth in relation to the carbon and nitrogen metabolic pathways under different photoperiod conditions.

2 Materials and Methods

2.1 Wastewater Medium Preparation

Prior to the experiment, a synthetic wastewater medium simulating a high strength municipal wastewater was prepared based on a modified wastewater composition

from Leong et al. [10] yielding concentrations of chemical oxygen demand, COD and ammonium-nitrogen, $\text{NH}_4^+\text{-N}$ at 145 mg/L and 48 mg/L, respectively. The following were compositions (mg/L) for the wastewater: sucrose (109), $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (10), CaCl_2 (42), $(\text{NH}_4)_2\text{SO}_4$ (226), K_2HPO_4 (180), KH_2PO_4 (35) and MgSO_4 (49).

2.2 *Chlorella Vulgaris Cultivation Using Synthetic Wastewater Medium*

Samples of the freshwater microalgae, *Chlorella vulgaris* was obtained from the culture collections belonging to the Centre for Biofuel and Biochemical Research (CBBR), Universiti Teknologi PETRONAS. The microalgae, *Chlorella vulgaris* was chosen in the study due to its ability to populate ponds naturally combined with its adaptability in tolerating high nutrient loadings in sewage effluents [11]. Stock culture of the microalgae, *Chlorella vulgaris* (500 mL) was inoculated in a 5-L laboratory clear glass bottle containing approximately 4.5 L of the prepared synthetic wastewater medium. The photobioreactor was continuously aerated with compressed air and illuminated with a white light-emitting diode (LED) light at the light intensity of 1200 lx. The initial cultivation pH was adjusted to 7.1 ± 0.1 .

2.3 *Photoperiod Experimental Design*

The experimental setup consisted of culturing the microalgae, *Chlorella vulgaris* acclimated to the synthetic wastewater medium under five different photoperiod design settings in 500 mL Erlenmeyer flasks as functional bioreactors. Approximately 50 mL of the *Chlorella vulgaris* inoculum was introduced in each of the bioreactor containing 450 mL of the synthetic wastewater medium. The different photoperiod designs in the study with varying light:dark cycle (h:h) included 24:0 (autotrophic), 16:8, 12:12, 8:16 and 0:24 (heterotrophic) designated as BR-24L, BR-16L, BR-12L and BR-8L and BR-0L, respectively (Table 1). All the bioreactors except for the BR24-L (autotrophic) were supplemented with the additional organic

Table 1 Designated bioreactors under varying photoperiod cultivation conditions

Bioreactor	Photoperiod	
	Light (h)	Dark (h)
BR-24L (Autotrophic)	24	0
BR-16L	16	8
BR-12L	12	12
BR-8L	8	16
BR-0L (Heterotrophic)	0	24

carbon source (sucrose) from the compositions of the synthetic wastewater medium. Each of the bioreactor was continuously aerated with compressed air and the cultivation pH was regulated and maintained at 7.0 ± 0.1 using either 1 N of H_2SO_4 or 1 N of NaOH throughout the study. The experimental study was carried out until the stationary growth phase was achieved and performed in duplicates. Samplings were executed every two days for analyzing the microalgal biomass, COD and NH_4^+-N concentrations.

2.4 Analytical Procedures

The microalgal biomass concentration was determined via spectrophotometry using a UV-VIS spectrophotometer (Shimadzu UV-2600). An analysis of optical density was measured at the wavelength of 688 nm and the microalgal biomass concentration, B (g/L) was calculated using Eq. 1.

$$B = 0.3557 \times OD_{688nm}, R^2 = 0.988 \quad (1)$$

The concentration of COD was determined using the closed reflux, titrimetric method: 5520 following the Standard Methods for the Examination of Water and Wastewater [12]. The concentrations of NH_4^+-N species were also analysed following the Standard Methods [12], namely, the titrimetric method: 4500-NH₃.


3 Results and Discussion

3.1 Microalgal Biomass Growth

The growth patterns of *Chlorella vulgaris* were visualized in Fig. 1. The initial microalgal biomass in all bioreactors are relatively the same, which was around

Fig. 1 Microalgal biomass growth cultured under different photoperiods

Microalgal biomass concentration (g/L) against time (day)



0.2 g/L. The first 2 days were considered as the lag phase period of microalgae cultured in all bioreactors as demonstrated by a relatively low biomass growth. BR-24L experienced a sudden increase in biomass concentrations (days 2–8) before entering the stationary phase at the 10th day of cultivation. BR-16L experienced a logarithmic growth phase in the 4th–8th day of cultivation whereas BR-12L and BR-8L experienced logarithmic growth phase later in the 8th–14th day and 10th–14th day of cultivation respectively. A trend was observed at which longer photoperiod induces faster microalgal biomass growth. Meanwhile, no logarithmic growth phase was observed in BR-0L indicating that the growth of *Chlorella vulgaris* was inhibited in heterotrophic culture.


At the end of cultivation study (14th day), the final biomass yields of 1.115 ± 0.003 g/L, 1.130 ± 0.001 g/L, 1.087 ± 0.040 g/L, 1.065 ± 0.020 g/L, and 0.359 ± 0.009 g/L were achieved in BR-24L, BR-16L, BR-12L, BR8-L, and BR-0L respectively. Although the final biomasses of BR-24L, BR-16L, BR-12L and BR-8L are comparable, BR-16L attained the highest biomass yield which suggested that under cyclic dark/hour conditions, cell production of biomass is higher.

3.2 Carbon (COD) Removal

COD measurement was carried out in the present study to estimate the carbon removal efficiencies in terms of COD removal from the synthetic wastewater media by *Chlorella vulgaris* at different photoperiods. The amount of organic carbon (sucrose) added to the bioreactors in this study generating an initial COD value of 103.63 mg/L. Figure 2 depicts the removal of COD under different photoperiods.

Rapid removal of COD was observed where about 80% COD removal was attained in all bioreactors on the second day of cultivation where similar COD removal patterns were noticed stipulating that there was no significant relationship between COD removal and cultivation of *Chlorella vulgaris* at different photoperiods. At the end of cultivation, the final concentrations of COD were found to be 6.72 ± 1.33 , 8.64 ± 3.991 , 13.44 ± 2.661 , and 13.44 ± 2.661 mg/L in BR16-L, BR12-L, BR-8L

Fig. 2 COD removal by *Chlorella vulgaris* cultivated under various photoperiods



and BR-0L respectively. The almost complete removal of COD from microalgae culture suggested that complete biodegradable COD source was immediately being assimilated by *Chlorella vulgaris*.

3.3 Nitrogen Removal


Microalgae is known to be able to utilize nutrients in particular nitrogen and phosphorus for its growth [13]. Nitrogen could present in various form such as nitrite (NO_2^-), nitrate (NO_3^-), ammonium ($\text{NH}_4^+\text{-N}$), ammonia (NH_3), nitric acid (HNO_3) and nitrogen gas (N_2). In the present study, $\text{NH}_4^+\text{-N}$ species was introduced in the synthetic wastewater medium. The removal of $\text{NH}_4^+\text{-N}$ species efficiencies by *Chlorella vulgaris* cultured at different photoperiods is represented in Fig. 3.

From Fig. 3, high $\text{NH}_4^+\text{-N}$ removal is observed in BR-24L followed by BR-12L, BR-16L and BR-0L with final concentration of $0, 0.38 \pm 0.527, 3.15 \pm 0.873, 15.75 \pm 0.873$ and 30.87 ± 0.873 respectively. BR-24L exhibited the fastest $\text{NH}_4^+\text{-N}$ removal at which complete removal was observed at the 8th day of cultivation at which the microalgal growth has reached stationary growth phase. A similar trend was observed in BR-16L, BR-12L, and BR-8L where high removal of $\text{NH}_4^+\text{-N}$ species is achieved as the microalgal growth entering stationary growth phase. Contrarily, the $\text{NH}_4^+\text{-N}$ uptake by *Chlorella vulgaris* grown in BR-0L was impeded as no decline in $\text{NH}_4^+\text{-N}$ concentrations is seen after second day of cultivation which assumed that $\text{NH}_4^+\text{-N}$ was not assimilated by microalgae.

It is interesting to note that, although the intake of nitrogen was hindered in BR-0L, a prominent decrease in COD is seen in the second day of cultivation. An assumption was made that a complete consumption of the organic carbon source i.e. sucrose, occurred during the second day of cultivation. The notorious decrease in COD values caused by the complete degradation of organic carbon sources by the microalgae which explained the stagnant growth of microalgae in BR-0L (heterotrophic) even after 14 days of cultivation. The growth of microalgae grown under total dark condition is highly dependent on the presence of organic carbon source. Once the organic

Fig. 3 Ammonium-nitrogen ($\text{NH}_4^+\text{-N}$) removal by *Chlorella vulgaris* cultivated under various photoperiods

Ammonium-Nitrogen concentration (mg/L) against time (day)



carbon source was used up, the microalgal growth will be ceased as there is no source of energy left to support the cell metabolism for microalgal growth. The reasoning is further supported by the intake of $\text{NH}_4^+\text{-N}$ species was prohibited as the assimilation of nitrogen into microalgal cell is largely associated with the availability of organic carbon source that serves as energy to convert $\text{NH}_4^+\text{-N}$ into biodegradable nitrogen for microalgae assimilation [14]. Nonetheless, other bioreactors continue to demonstrate an increase in biomass growth and reduction in nitrogen concentration although organic compound was fully consumed. It can be inferred that after second day of cultivation, the growth of microalgae in other bioreactors were solely rely on the light energy via the photosynthetic pathway. It is also worth noting that higher $\text{NH}_4^+\text{-N}$ removal efficiencies were achieved at longer photoperiods which is in tandem with the trend observed in the microalgal biomass growth.

4 Conclusion

This study aimed at finding the optimum photoperiods in enhancing biomass and lipid productions by investigating the effect of photoperiods on the biomass production, carbon and nutrient uptakes and lipid yields. Five different photoperiods were chosen to study in this study. Higher microalgal biomass yield accompanied by higher nitrogen removal by *Chlorella vulgaris* were observed under longer light photoperiod cycles. However, there were no remarkable differences observed in the light cycles of 24 h, 16 h and 12 h in the photoperiod study. Therefore, the 12 h photoperiod duration is deemed to be the most ideal cultivation condition considering lower energy consumption for illumination viewing from an economic perspective. It can be deduced that the concentration of organic carbon source i.e. sucrose, introduced in the microalgal growth culture is relatively low leading to carbon deficit in the microalgal consortium. Dark/light cycle can counterbalance the shortage of organic carbon source by utilizing carbon dioxide as an inorganic carbon source in the presence of light for photosynthetic metabolic grow. As a result, high biomass growth was achieved under the 16 h, 12 h and 8 h light cycles of photoperiod whereas the growth of the heterotrophic condition (0 h) was totally inhibited.

Acknowledgements The financial supports from the Ministry of Higher Education Malaysia via Fundamental Research Grant Scheme (FRGS) with the cost center of 015MA0-110 (FRGS/1/2020/TK0/UTP/02/20), HICoE-Center for Biofuel and Biochemical Research with the cost center of 015MA0-052, Research Collaboration Grant UTP-UMP-UMT-UCTS with the cost center of 015MD0-019 and Yayasan Universiti Teknologi PETRONAS via YUTP-FRG with the cost center of 015LC0-126 are gratefully acknowledged.

References

1. Rashid, N., Rehman, M.S.U., Sadiq, M., Mahmood, T., Han, J.I.: Current status, issues and developments in microalgae derived biodiesel production. *Renew. Sustain. Energy Rev.* **40**, 760–778 (2014)
2. Khan, S.A., Rashmi, Hussain, M.Z., Prasad, S., Banerjee, U.C.: Prospects of biodiesel production from microalgae in India. *Renew. Sustain. Energy Rev.* **13**(9), 2361–2372 (2009)
3. Salama, E.S., Kurade, M.B., Abou-Shanab, R.A., El-Dalatony, M.M., Yang, I.S., Min, B., Jeon, B.H.: Recent progress in microalgal biomass production coupled with wastewater treatment for biofuel generation. *Renew. Sustain. Energy Rev.* **79**, 1189–1211 (2017)
4. Khoo, K.S., Chew, K.W., Yew, G.Y., Leong, W.H., Chai, Y.H., Show, P.L., Chen, W.H.: Recent advances in downstream processing of microalgae lipid recovery for biofuel production. *Bioresour. Technol.* **304**, 122996 (2020)
5. Vo, H.N.P., Ngo, H.H., Guo, W., Chang, S.W., Nguyen, D.D., Chen, Z., Wang, X.C., Chen, R., Zhang, X.: Microalgae for saline wastewater treatment: a critical review. *Crit. Rev. Environ. Sci. Technol.* **50**(12), 1224–1265 (2020)
6. Lam, M.K., Lee, K.T.: Microalgae biofuels: a critical review of issues, problems and the way forward. *Biotechnol. Adv.* **30**(3), 673–690 (2012)
7. da Silva Ferreira, V., Sant’Anna, C.: The effect of physicochemical conditions and nutrient sources on maximizing the growth and lipid productivity of green microalgae. *Phycol. Res.* **65**(1), 3–13 (2017)
8. Fields, M.W., Hise, A., Lohman, E.J., Bell, T., Gardner, R.D., Corredor, L., Moll, K., Peyton, B.M., Characklis, G.W., Gerlach, R.: Sources and resources: importance of nutrients, resource allocation, and ecology in microalgal cultivation for lipid accumulation. *Appl. Microbiol. Biotechnol.* **98**(11), 4805–4816 (2014)
9. Leong, W.H., Lim, J.W., Lam, M.K., Uemura, Y., Ho, Y.C.: Third generation biofuels: a nutritional perspective in enhancing microbial lipid production. *Renew. Sustain. Energy Rev.* **91**, 950–961 (2018)
10. Leong, W.H., Kiatkittipong, K., Kiatkittipong, W., Cheng, Y.W., Lam, M.K., Shamsuddin, R., Mohamad, M., Lim, J.W.: Comparative performances of microalgal-bacterial co-cultivation to bioremediate synthetic and municipal wastewaters whilst producing biodiesel sustainably. *Processes* **8**(11), 1427 (2020)
11. Leong, W.H., Lim, J.W., Lam, M.K., Uemura, Y., Ho, C.D., Ho, Y.C.: Co-cultivation of activated sludge and microalgae for the simultaneous enhancements of nitrogen-rich wastewater bioremediation and lipid production. *J. Taiwan Inst. Chem. Eng.* **87**, 216–224 (2018)
12. APHA, AWWA, WEP: Standard Methods for the Examination of Water and Wastewater, 22nd ed. American Public Health Association, Washington, DC (2012)
13. Perez-Garcia, O., Bashan, Y.: Microalgal heterotrophic and mixotrophic culturing for bio-refining: from metabolic routes to techno-economics. In: *Algal Biorefineries*, pp. 61–131. Springer, Cham (2015)
14. Perez-Garcia, O., Escalante, F.M., de-Bashan, L.E., Bashan, Y.: Heterotrophic cultures of microalgae: metabolism and potential products. *Water Res.* **45**(1), 11–36 (2011)

