

Krishna Prasad Vadrevu · Toshimasa Ohara ·
Chris Justice
Editors

Vegetation Fires and Pollution in Asia

 Springer

Editors

Krishna Prasad Vadrevu
NASA Marshall Space Flight Center
Huntsville, AL, USA

Toshimasa Ohara
Center for Environmental Science
in Saitama
Kazo, Saitama, Japan

Chris Justice
University of Maryland
College Park, MD, USA

ISBN 978-3-031-29915-5 ISBN 978-3-031-29916-2 (eBook)
<https://doi.org/10.1007/978-3-031-29916-2>

© The Editor(s) (if applicable) and The Author(s), under exclusive license to Springer Nature Switzerland AG 2023

All rights are solely and exclusively licensed by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

Foreword



It is my pleasure to write a Foreword to the book on *Vegetation Fires and Air Pollution in Asia*, edited by Drs. Krishna Prasad Vadrevu, Toshimasa Ohara, and Chris Justice. I wrote the forewords to their earlier books: (1) *Land-Atmosphere Interactions in South/Southeast Asia* (2018) and (2) *Biomass Burning in South and Southeast Asia* (2022) that drew wide attention in the research community. Wildfires are among the most serious disasters over the globe and in South and Southeast Asian countries, in particular. Transboundary pollution from vegetation fires is a recurrent environmental problem in Asian countries. Addressing fire-related impacts on ecosystems and the atmosphere requires robust information on their spatiotemporal characteristics, including impact assessment and uncertainties. Remote sensing and geospatial technologies offer unique opportunities for mapping and monitoring fires and emissions estimation, and impacts assessment needed for land management and air pollution control.

The current book is a compilation of articles on various fire-related topics from renowned researchers in the USA and South/Southeast Asia. The contributions in

scientists, this book is a significant addition to the literature on fire research and applications and biomass burning in South/Southeast Asia. I am impressed with the book's content and the rich contributions from regional scientists. The book is a timely contribution to NASA LCLUC SARI science and reflects the current status in this research field.

The editors of this book are highly active and reputed researchers who have published multiple books earlier. I commend all the team members, particularly the Chief Editor, Dr. Krishna Prasad Vadrevu, SARI Lead, on accomplishing another big task of compiling rich material. I am confident it will motivate new, exciting research on fire and emissions and trigger innovative ideas. I welcome more such contributions from the US and regional scientists and wish you all good reading.

Dr. Garik Gutman
Land-Cover/Land-Use Change
Program Manager
NASA Headquarters
Washington DC, USA

Reference

Gutman, G., Csiszar, I., and P. Romanov. 2000. Using NOAA/AVHRR products to monitor El Nino impacts: Focus on Indonesia in 1997–98. *Bulletin of the American Meteorological Society* 81: 1189–1205.

Contents

Vegetation Fires and Biomass Burning in South/Southeast Asia—An Overview	1
Krishna Prasad Vadrevu, Toshimasa Ohara, and Chris Justice	
Mapping, Monitoring, and Modeling of Vegetation Fires	
Wildfire Monitoring Using Infrared Bands and Spatial Resolution Effects	21
Wei Zheng, Jie Chen, Jinlong Fan, Yajun Li, and Cheng Liu	
Status and Drivers of Forest Fires in Myanmar	35
Sumalika Biswas and Krishna Prasad Vadrevu	
Vegetation Fires and Entropy Variations in Myanmar	53
Krishna Prasad Vadrevu, Pranith Salikineedi, Aditya Eaturu, and Sumalika Biswas	
Crop Residue Burning and Forest Fire Emissions in Nepal	71
Bhupendra Das, Siva Praveen Puppala, Bijaya Maharjan, Krishna B. Bhujel, Ajay Mathema, Dhurba Neupane, and Rejina M. Byanju	
Firewood Burning Dynamics by the Sri Lankan Households: Trends, Patterns, and Implications	85
Asankha Pallegedara and Ajantha Sisira Kumara	
Burnt Area Signal Variations in Agriculture and Forested Landscapes of India—A Case Study Using Sentinel-1A/B Synthetic Aperture Radar	99
Krishna Prasad Vadrevu, Aditya Eaturu, and Sumalika Biswas	

Application of Interferometry SAR for Monitoring of Peatland Area—Case Studies in Indonesia 119
 Yessy Arvelyna, Prayoto Pranoto, Keiko Ishi, and Krishna Prasad Vadrevu

Active Fire Monitoring of Thailand and Upper ASEAN by Earth Observation Data: Benefits, Lessons Learned, and What Still Needs to Be Known 139
 Veerachai Tanpipat, Jessica L. McCarty, Diane Davies, Wilfrid Schroeder, and Chris Elvidge

Detecting Vegetation Regrowth After Fires in Small Watershed Settings Using Remotely Sensed Data and Local Community Participation Approach 155
 Thaworn Onpraphai, Attachai Jintrawet, Angkana Somsak, Suprapat Khuenjai, Pong Loungmoon, Bounthanh Keoboulapha, and Jun Fan

Long-Term Spatiotemporal Distribution of Fire Over Maritime Continent and Their Responses to Climate Anomalies 173
 Aulia Nisa’ul Khoir, Maggie Chel Gee Ooi, and Nur Nazmi Liyana Binti Mohd Napi

Vegetation Fires in Laos—An Overview 187
 Krishna Prasad Vadrevu, Chittana Phompila, and Aditya Eaturu

Vegetation Fires, Fire Radiative Power, and Intermediate Fire Occurrence-Intensity (IFOI) Hypothesis Testing in Myanmar, Laos, and Cambodia 209
 Krishna Prasad Vadrevu, Aditya Eaturu, Thav Sopheak, Chittana Phompila, and Sumalika Biswas

Analyzing Fire Behavior and Calibrating a Fire Growth Model in a Seasonally Dry Tropical Forest Area 233
 Gernot Ruecker, Veerachai Tanpipat, and Kobsak Wanthongchai

Greenhouse Gas Emissions and Air Pollution

Spatiotemporally Resolved Pollutant Emissions from Biomass Burning in Asia 255
 Rui Xiong, Yatai Men, and Guofeng Shen

Twenty-Year (2000–2019) Variations of Aerosol Optical Depth Over Asia in Relation to Anthropogenic and Biomass Burning Emissions 269
 Syuichi Itahashi, Junichi Kurokawa, and Toshimasa Ohara

Light Absorption Properties of Biomass Burning Emissions in Bangladesh: Current State of Knowledge	283
Shahid Uz Zaman, Md Safiqul Islam, Shatabdi Roy, Farah Jeba, and Abdus Salam	
Remote Sensing of Greenhouse Gases and Aerosols from Agricultural Residue Burning Over Pakistan	299
Salman Tariq, Hasan Nawaz, Zia Ul-Haq, and Usman Mehmood	
A Comparative Study of Energy, Emissions, and Economic Efficiency of Various Cookstoves in Nepal	317
Narayan P. Adhikari, Prajwal R. Shakya, Shubha Laxmi Shrestha, and Suyesh Prajapati	
Estimation of Ultrafine Particulate Matter Emissions from Biomass Burning Using Satellite Imaging and Burn Severity	339
Perapong Tekasakul, Narissara Nuthammachot, Rachane Malinee, John Morris, and Racha Dejchanchaiwong	
Characteristics of Transboundary Haze and General Aerosol Over Pulau Pinang, Malaysia	359
Lim Hwee San, Brent N. Holben, Ezekiel Kaura Makama, and Mohamad Farid Izzat Bin Zahari	
Measurements of Atmospheric Carbon Dioxide Emissions from Fire-Prone Peatlands in Central Kalimantan, Indonesia, Using Ground-Based Instruments	385
Masafumi Ohashi, Windy Iriana, Osamu Kozan, Masahiro Kawasaki, and Kenichi Tonokura	
Air Pollution Caused by Peatland Fires in Central Kalimantan	401
Hiroshi Hayasaka and Aswin Usup	
Chemical Speciation of PM₁₀ Emissions from Peat Burning Emission in Central Kalimantan, Indonesia	417
Puji Lestari, Isna Utami, Febri Juwita, Rajasekhar Balasubramanian, and Jeffrey S. Reid	
GHG Emissions' Estimation from Peatland Fires in Indonesia—Review and Importance of Combustion Factor	433
Bambang Hero Saharjo	
Forest Fire Emissions in Equatorial Asia and Their Recent Delay Anomaly in the Dry Season	447
Ida Bagus Mandhara Brasika	

Air Pollution Modeling and Decision Support Systems

- Impact of Vegetation Fires on Regional Aerosol Black Carbon Over South and East Asia** 465
Yogesh Kant, Aryan Natwariya, Debashis Mitra, and Prakash Chauhan
- Detection and Modeling of South Asian Biomass Burning Aerosols from Both Macro- and Micro-perspectives** 483
Shuaiyi Shi, Tianhai Cheng, Yu Wu, Xingfa Gu, Xiaoyang Li, Siheng Wang, and Yuyang Wang
- Remote Sensing of Agricultural Biomass Burning Aerosols, Gaseous Compounds, Long-Distance Transport, and Impact on Air Quality** 499
Yonghua Wu, Yong Han, and Fred Moshary
- Agricultural Fires in Northeast China: Characteristics, Impacts, and Challenges** 517
Jiumeng Liu and Yuan Cheng
- Air Pollution Modeling in Southeast Asia—An Overview** 531
Teerachai Amnuaylojaroen
- Trace Gases and Air Quality in Northwestern Vietnam During Recurrent Biomass Burning on the Indochina Peninsula Since 2014—Field Observations and Atmospheric Simulations** 545
Simone M. Pieber, Stephan Henne, Nhat Anh Nguyen, Dac-Loc Nguyen, and Martin Steinbacher
- Southeast Asian Transboundary Haze in the Southern Philippines, 2019 and Meteorological Drivers** 559
Krishna E. Santos, Mylene G. Cayetano, and Prisco D. Nilo
- An Operational Fire Danger Rating System for Thailand and Lower Mekong Region: Development, Utilization, and Experiences** 575
Veerachai Tanpipat, Kasemsan Manomaiphiboon, Robert D. Field, William J. deGroot, Prayoonyong Nhuchaiya, Narin Jaronrattanapak, Chatchaya Buaniam, and Jittisak Yodcum
- Fires Hotspot Forecasting in Indonesia Using Long Short-Term Memory Algorithm and MODIS Datasets** 589
Evizal Abdul Kadir, Hsiang Tsung Kung, Arbi Haza Nasution, Hanita Daud, Amal Abdullah AlMansour, Mahmud Othman, and Sri Listia Rosa



Dr. Chris Justice is a professor at the Department of Geographical Sciences, University of Maryland, College Park, USA. He has 40 years of research experience. His current research is on land-cover and land-use change and global agricultural monitoring using remote sensing. He is an authority on satellite remote sensing of fires. He serves as Project Scientist for the NASA LCLUC Program, the Land Discipline Lead for the NASA MODIS and the Suomi-NPP VIIRS Science Team. He is Co-Chair of the GEO Global Agricultural Monitoring initiative (GEOGLAM), Chief Scientist for NASA HARVEST, and Chair of the international Global Observations of Forest and Land-Use Dynamics (GOFC-GOLD) program.

Contributors

Narayan P. Adhikari Alternative Energy Promotion Center, Kathmandu, Nepal

Amal Abdullah AlMansour Department of Computer Science, King Abdul Aziz University, Jeddah, Saudi Arabia

Teerachai Amnuaylojaroen School of Energy and Environment, University of Phayao, Phayao, Thailand

Yessy Arvelyna Remote Sensing Technology Center of Japan, Tokyo, Japan

Rajasekhar Balasubramanian Department of Civil and Environmental Engineering, Faculty of Engineering, National University of Singapore, Singapore, Singapore

Krishna B. Bhujel Nepal Energy and Environment Development Services (NEEDS), Kathmandu, Nepal

Sumalika Biswas University of California, Los Angeles, USA

Ida Bagus Mandhara Brasika Department of Marine Science, Udayana University, Bali, Indonesia;
Department of Mathematics and Statistics, The University of Exeter, Exeter, UK

Chatchaya Buaniam Department of National Parks, Wildlife, and Plant Conservation, Ministry of Natural Resources and Environment, Bangkok, Thailand

Rejina M. Byanju Central Department of Environmental Science, Tribhuvan University, Kirtipur, Nepal

Mylene G. Cayetano Institute of Environmental Science and Meteorology, University of the Philippines-Diliman, Quezon City, Philippines

Fires Hotspot Forecasting in Indonesia Using Long Short-Term Memory Algorithm and MODIS Datasets



Evizal Abdul Kadir, Hsiang Tsung Kung, Arbi Haza Nasution, Hanita Daud, Amal Abdullah AlMansour, Mahmud Othman, and Sri Listia Rosa

Abstract Vegetation fires are most common in South and Southeast Asian countries, including Indonesia. In addition to anthropogenic causes, climate change in the form of droughts is the biggest driver of fires in Indonesia. In particular, the peatlands in Indonesia are highly vulnerable to droughts with recurrent fires. In this study, we used a long short-term memory (LSTM) algorithm to predict the fire hotspots based on the 2010 to 2021 fire data. More than 700,000 fire hotspots from 2010 to 2021 have been collected and used as a training dataset to forecast fires for the year 2022. The LSTM algorithm successfully predicted 2022 fires with the minimum root mean squared error and high accuracy. Furthermore, the results of the 2022 prediction year matched the previous year's fire data seasonally, with increasing fires from August to November. The study highlights the potential use of the LSTM algorithm for forecasting fires in Indonesia.

Keywords Fires hotspot · Forecasting · Indonesia · LSTM · MODIS

1 Introduction

Fires are one of the biggest natural threats to forests, woodlands, and grasslands in many countries, including Indonesia (Albar et al. 2018; Akther and Hassan, 2011; Goldammer 2012; Hayasaka et al. 2014; Petropoulos et al. 2013; Justice et al. 2015; Kadir et al. 2019, 2020, 2021). In several South/Southeast Asian countries, fire is used to clear the forests for agriculture through slash and burn (Albar et al. 2018;

E. A. Kadir (✉) · A. H. Nasution · S. L. Rosa
Department of Informatics Engineering, Universitas Islam Riau, Pekanbaru 28284, Indonesia
e-mail: evizal@eng.uir.ac.id

E. A. Kadir · H. T. Kung
Department of Computer Science, Harvard University, Cambridge, MA 02134, USA

H. Daud · M. Othman
Department of Applied Mathematics, Universiti Teknologi Petronas, 86400 Perak, Malaysia

A. A. AlMansour
Department of Computer Science, King Abdul Aziz University, Jeddah 80200, Saudi Arabia

© The Author(s), under exclusive license to Springer Nature Switzerland AG 2023

589

K. Vadrevu et al. (eds.), *Vegetation Fires and Pollution in Asia*,
https://doi.org/10.1007/978-3-031-29916-2_35

Badarinath et al. 2007; 2008, 2009, 2015b; Badarinath and Prasad 2011; Biswas et al. 2015a); Biswas et al. 2021; Kant et al. 2000; Lasko and Vadrevu 2018; Petropoulos et al. 2013; Prasad et al. 2001a,b; Prasad et al. 2002a, 2002b;2003; 2004; Prasad and Badarinath 2004; Prasad et al. 2005; Prasad and Badarinth 2006; Vadrevu 2008; 2021a,b; Wooster et al. 2021; (Biswas et al. 2015a,b; Prasad et al. 2001a, b, 2002a, b;) agricultural residues after crop harvest to clear the land for the next crop (Lasko et al. 2017; 2018a,b; 2021; Vadrevu and Lasko 2015), to clear the forested lands for plantations (Albar et al. 2018), promoting the growth of grass in pasture lands for cattle (Thapa et al. 2022), etc., in addition to intentional or accidental human activities. While most of these fires are anthropogenic, the drivers of fires can also be natural such as lightning and extreme and prolonged drought conditions. Especially in tropical regions, there are usually two alternating rainy and dry seasons, and forests and grassland fires are highly vulnerable to fires during the dry season. Indonesia is one of the tropical countries with major fire issues, especially in Kalimantan and Sumatra Islands with recurrent fires (Hayasaka et al. 2014). Regardless of the ignition source, in forested areas, the fires can spread rapidly and become uncontrollable due to the local meteorological and environmental conditions. Further, fires are a major important source of air pollution which results in the release of greenhouse gas emissions and aerosols (Ito and Penner 2005; Gupta et al. 2001; Lasko and Vadrevu 2018; Vadrevu and Badarinath 2009; Vadrevu and Justice 2011; Kharol et al. 2012; Vadrevu and Lasko 2015; Vadrevu 2015; Vadrevu et al. 2008; 2013;). The smoke particles released from fires can interact with the cloud droplets and alter Earth's radiation budget (Martins and Dias 2009). The GHG emissions from biomass burning represent the largest source of inter-annual variability, in particular, CO₂ fluxes (Szopa et al. 2007; Kant et al. 2000;). Biomass burning is estimated to contribute to 7600 ± 359 Tg CO₂eq year⁻¹ (FAOSTAT 2020). In addition, biomass burning has been shown to influence various land-atmospheric interactions at different scales, such as vegetation transpiration, soil erosion, albedo (Crutzen and Andreae 1990). Smoke-borne aerosols from fires disrupt normal hydrological processes and reduce rainfall, potentially contributing to regional drought. In addition to these effects on Earth's radiation, atmosphere, climate, and ecosystems, the pollutants released from the fires (Vadrevu et al. 2014a,b, 2017, 2018 2019) can impact health resulting in asthma, acute respiratory illness, eye irritation, cardiovascular mortality, thrombosis, etc. (Sigsgaard et al. 2015). Thus, fires can become a disaster for humans and the environment due to their severity and intensity. Considering these effects, mapping and monitoring of fires, including forecasting, can not only help in understanding land-atmospheric interactions useful for climate change studies but also protecting human lives, ecosystems, and related functions (Goldammer 2012; Eaturu and Vadrevu 2021; Vadrevu and Justice 2011; Vadrevu et al. 2020; 2021a,b, 2022a; b).

Several techniques have been proposed to forecast fires, such as fire danger indices combining climate data with site characteristics and fire data records (Akhter and Hassan 2011; Vadrevu et al. 2021a, b). In addition, multiple machine learning algorithms were also used to characterize fire patterns and predict fires. For most algorithms, previous fire data is essential for calibration and prediction (Liang et al. 2019;

Omar et al. 2021; Lamjiak et al. 2021; Abdul Kadir et al. 2022; Mohan et al. 2021). These studies considered both the climate and environmental factors in predicting the fires. Including meteorological factors in the prediction of fires is important as they can drive accuracy. A comprehensive data analysis of fire hotspot occurrences, their fire size, intensity, and how they can potentially spread into new areas, including forecasting methods, were given in earlier works (Khabarov et al. 2008; Han et al. 2019; Kadir et al. 2019; Kukuk and Kilimci 2021; Prapas et al. 2021). Recently, deep learning algorithms are gaining popularity in various fields, such as pattern recognition, including forecasting (Benzekri et al., 2020). In this study, we use the popular long short-term memory (LSTM) algorithm to forecast fires in Indonesia for 2022. We used the fire spots data derived from the Moderate Resolution Imaging Spectroradiometer (MODIS) from 2010 to 2021 and tested the algorithm's robustness in predicting the fires for 2022.

2 Datasets and Methodology

We used the NASA MODIS fire hotspots data from 2010–2021 for our study. Table 1 shows the sample fires dataset for Indonesia. The data has been normalized and grouped into a single date of fire occurrence. The data has been split into training and testing for fire forecasting. In the field of deep learning, the LSTM algorithm is an artificial recurrent neural network (RNN) architecture and was first introduced by Hochreiter and Schmidhuber (1997). LSTM is a special model of RNN that capable of learning in long-term dependencies and remembering information for prolonged periods as a default. Figure 1 shows the RNN-LSTM model's architecture, consisting of several main blocks called cells with input, output, and forget gates. The sigmoid activation function classifies the values in probabilities for the two predefined classes in the dense output layer.

The LSTM model can be explained as short-term memory, which acts when the information is being acquired, retains for a few seconds, and then destines it to be kept for more extended periods or discards it. Long-term memory permanently retains information, allowing its recovery or recall. It contains all our autobiographical data and all our knowledge. LSTM model can handle the problem with long-term dependencies of RNN in which the RNN algorithm cannot do in the prediction of the information stored in the long-term memory but can give more accurate prediction from the recent information. LSTM can use by default to retain the data for a long-term period. The algorithm can predict, process, and classify based on time series data (Le et al. 2019). The LSTM model has an incredible way of forecasting and works well in time series data. Furthermore, this model can organize in the form of a chain structure and has four interacting layers with a unique method of communication in data processing. Figure 2 shows an analysis block diagram of how the forecasting process of the fire hotspot is done in our study.

Table 1 Detail of fires hotspot dataset from year 2010 to 2021 (NASA 2021)

	Latitude	Longitude	Brightness	Scan	Track	acq_data	acq_time	Satellite	Instrument	Confidence	Version	Bright_β1	Frp	Daynight	Type
0	0.02110	116.87390	315.30	1.10	1.10	2010-01-01	251	terra	MODIS	42	6.2	295.60	8.70	D	0.0
1	0.48080	116.08060	312.30	1.00	1.00	2010-01-01	251	terra	MODIS	66	6.2	295.00	6.90	D	0.0
11	2.15090	117.49680	320.60	1.00	1.00	2010-01-01	550	Aqua	MODIS	0	6.2	297.50	10.60	D	0.0
10	8.10890	118.07430	319.30	1.00	1.00	2010-01-01	547	Aqua	MODIS	0	6.2	300.80	9.10	D	0.0
8	8.15960	117.58570	319.60	1.00	1.00	2010-01-01	547	Aqua	MODIS	43	6.2	297.70	9.10	D	0.0
...
14208	7.22331	110.42920	316.80	1.00	1.00	2021-12-31	300	terra	MODIS	67	6.1NRT	293.096	8.56	D	NaN
14209	6.96059	110.45844	318.84	1.00	1.00	2021-12-31	300	terra	MODIS	55	6.1NRT	291.66	6.42	D	NaN
14210	5.80178	139.61118	309.54	1.00	1.00	2021-12-31	418	Aqua	MODIS	65	6.1NRT	283.90	5.50	D	NaN
14211	4.51654	136.84802	313.68	1.21	1.09	2021-12-31	418	Aqua	MODIS	56	6.1NRT	291.74	7.84	D	NaN
14213	4.54666	136.77507	309.21	1.22	1.10	2021-12-31	418	Aqua	MODIS	52	6.1NRT	287.79	5.59	D	NaN

703116 rows × 15 columns

Fig. 1 Structure of RNN-LSTM algorithm





Fig. 2 Approach followed for fire forecasting using LSTM



The first step in data processing in forecasting is to construct an LSTM network model to identify the inputs and eliminate the information that is not necessary for the cell structure of LSTM (Fig. 1). The process of identifying and excluding data is governed by the sigmoid function, which takes the output of the last LSTM unit h_{t-1} at time $t - 1$ and the current input X_t at time t . Additionally, the sigmoid function determines which part from the old output should be eliminated. This gate is called the forget gate f_t ; where f is a vector with values ranging from 0 to 1, corresponding to each number in the cell state, C_{t-1} . . Our collected data had more than 700,000 fire hotspots within 12 years and, after normalization, became 4365 datasets of fires grouped in each day. The data was divided into training and testing data (Fig. 2). The optimization process was followed to evaluate results, increase the performance and enhance accuracy to minimize the error and final forecasting. The LSTM cell with sigmoid function W_f and b_f are the weight matrices and bias, respectively, of the forget gate. This step decides and stores the input data from the new information X_t




Fig. 3 Internal LSTM model process

in the cell state and updates the cell state. Then, the sigmoid layer decides whether the new data should be updated or ignored (0 or 1), and the tanh h function gives weight to the values which is passed by deciding their level of importance (1 to 1). The two values are multiplied to update the new cell state. This new memory is then added to the old memory C_{t-1} resulting in C_t . Figure 3 depicts how the neuron process of the LSTM model works (Chen et al., 2021).

The next step is C_{t-1} and C_t are the cell states in the LSTM cell at time C_{t-1} and t while W and b are the weight matrices and bias of the cell state. In the last step, the value of h_t is based on the output cell state o_t , a sigmoid layer decides which parts of the cell state make it to the output. Next, the output of the sigmoid gate o_t is multiplied by the new values created by the \tanh layer from the cell state C_t , with a value ranging between 1 and 1. Finally, the performance of the fire forecasting was done using the root mean square error (RMSE) with the prediction and actual data values using the below equation (1).

$$RMSE = \sqrt{\frac{\sum_{i=1}^n (X_i - \hat{X}_i)^2}{n}} \tag{1}$$

In the equation, X_i and X'_i are the actual fires hotspot data compared to forecasting fires data at the time t ; X_i is the mean of actual values fires data and n is the total number of data. The smaller the RMSE values, the better the prediction.

3 Results and Discussion

Our fire dataset consisted of several parameters such as coordinate or location of fire occurrence, date and time, confidence level (probability of becoming a big fire and spreading out), brightness, day or nighttime, etc. (Table 1). In addition, we specifically used parameters that have a major impact and are essential to forecasting, which includes coordinates (latitude and longitude), acquisition date (*acq_date*), and confidence level. Figure 4a shows the mapping of fire hotspot distribution in Indonesia for 2021 and Fig. 4b for 2020. The fire hotspots were classified into five confidence levels, starting with the lowest from 0, low impact, and less potential to spread till 100, with high impact and high probability spread potential to become a big fire. The five-level classifications with confidence levels are shown in different colors (0–20 blue dot; level 21–40 green; 41–60 yellow; 61–80 as orange and 81–100 red with the highest).

The month-wise fire distribution is shown in Fig. 5a, b for the years 2021 and 2020, respectively. Classification based on confidence level and the distribution matched well with the total number of hotspots. Mostly, the map showed yellow and orange colors with confidence levels varying from 41–60 and 61–80, respectively. While red color is the highest potential of fire hotspots spread, they showed less in number in the predicted map.

Results from the LSTM suggested a similar pattern and number of daily fire hotspot incidents, with a maximum of 600 to 700 from the September to November dry or summer season. The daily average number of hotspots is 87. Although this number is insignificant for the entire of Indonesia, the number might increase drastically due to the prevailing weather and other fire-favorable factors. Another issue is the type of land that gets affected due to fires. For example, the Sumatra and Kalimantan Islands peatlands are easily ignited when dry land and fires are difficult to control. The LSTM algorithm for forecasting fire hotspots in Indonesia has been tested preliminarily to the 2121 data before 2022. Figure 6 compares actual fire hotspot data and forecasting results for the year 2021; the results showed a good agreement between the graphs. Preliminary forecasting suggested an RMSE error of 4.56%. We then fine-tuned the LSTM forecasting algorithm for 2022 by training more than 4000 datasets using the filtered data from 2010 to 2021; in essence, 30% of the total data was used for training and the rest 70% for testing.

Figure 7 shows a good agreement and similar normal distribution patterns for all the years, i.e., 2020, 2021, and 2022. The high occurrence of fire hotspots detected in the early part of the year, i.e., March, and lesser in the middle of the year, then increasing from September to November, is a typical pattern reflected in the figures. The spikes during the few days in late September are attributed to seasonal fires in Sumatra Island.

Overall, the LSTM RNN algorithm showed successful results with minimum error. The results of the 2022 prediction year matched the previous 2021-year fire data. Forecasting results in 2022 show good agreement and a similar pattern of fires with increasing fires from August to November. By comparing the predicted data




Fig. 4 Mapping of fires hotspots in Indonesia **a** year 2021 **b** year 2020

with the previous year’s data, we could achieve an accuracy of up to 95% with an RMSE error of 4.56%. More robust data is required on the local conditions to achieve further high accuracy at specific locations. Our future studies will focus on the same, i.e., collecting and analyzing the data at a much higher spatial resolution for different regions in Indonesia.

4 Conclusion

We demonstrated the long short-term memory (LSTM) algorithm’s potential in predicting and forecasting fire hotspots in Indonesia. A fire hotspots dataset from 2010 to 2021 obtained from the NASA MODIS data has been used to train and forecast fires for 2022. By comparing the predicted data with the previous year’s




Fig. 5 Distribution of fires hotspots in Indonesia for the year from January to December **a** year 2021 **b** year 2020




Fig. 6 Comparison of actual and fire forecasting data for year 2021




Fig. 7 Forecasting of fires hotspots in year 2022 and actual data of fires in year 2020–2021

data, we could achieve an accuracy of up to 95% with an RMSE error of 4.56%. The forecasted fire data patterns matched the previous year’s data in seasonality from January to December. It is noted that the number of hotspots increase by the end of each year due to the dry season in Indonesia.

Acknowledgements We thank the Ministry of Education, Culture, Research, and Technology of Indonesia for funding the research and American Indonesia Exchange Foundation (AMINEF) and the Fulbright fellowship. We also acknowledge Harvard University, Universiti Teknologi Petronas, Universitas Islam Riau, and King Abdul Aziz University for their research facilities.

References

- Abdul Kadir, E., S. Listia Rosa, A. Syukur, M. Othman, and H. Daud. 2022. Forest fire spreading and carbon concentration identification in tropical region Indonesia. *Alexandria Engineering Journal* 61: 1551–1561.
- Alkher, M.S., and Q.K. Hassan. 2011. Remote sensing-based assessment of fire danger conditions over boreal forest. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* 4: 992–999.
- Albar, I., I. Jaya, B.H. Saharjo, B. Kuncahyo, K.P. Vadrevu. (2018). Spatio-temporal analysis of land and forest fires in Indonesia using MODIS active fire dataset. In *Land-atmospheric research applications in South and Southeast Asia*, (pp. 105–127). Springer, Cham.
- Badarinath, K.V.S., and K.V. Prasad. 2011. Carbon dioxide emissions from forest biomass burning in India. *Global Environmental Research* 15: 45–52.
- Badarinath, K.V.S., S.K. Kharol, K.M. Latha, T.K. Chand, V.K. Prasad, A.N. Jyothsna, and K. Samatha. 2007. Multiyear ground-based and satellite observations of aerosol properties over a tropical urban area in India. *Atmospheric Science Letters* 8 (1): 7–13.
- Badarinath, K.V.S., S.K. Kharol, A.R. Sharma, and V.K. Prasad. 2009. Analysis of aerosol and carbon monoxide characteristics over Arabian Sea during crop residue burning period in the Indo-Gangetic Plains using multi-satellite remote sensing datasets. *Journal of Atmospheric and Solar-Terrestrial Physics* 71 (12): 1267–1276.
- Badarinath, K.V.S., S. Kumar Kharol, V. Krishna Prasad, A. Rani Sharma, E.U.B. Reddi, H.D. Kambezidis, and D.G. Kaskaoutis. 2008. Influence of natural and anthropogenic activities on UV Index variations—a study over tropical urban region using ground based observations and satellite data. *Journal of Atmospheric Chemistry* 59 (3): 219–236.
- Benzekri, W., A.E. Moussati, O. Moussaoui, and M. Berrajaa. 2020. Early forest fire detection system using wireless sensor network and deep learning. *International Journal of Advanced Computer Science and Applications (IJACSA)* 11: 496–503.
- Biswas, S., K.D. Lasko, and K.P. Vadrevu. 2015a. Fire disturbance in tropical forests of Myanmar—analysis using MODIS satellite datasets. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* 8 (5): 2273–2281.
- Biswas, S., K.P. Vadrevu, Z.M. Lwin, K. Lasko, and C.O. Justice. 2015b. Factors controlling vegetation fires in protected and non-protected areas of Myanmar. *PLoS ONE* 10 (4): e0124346.
- Biswas, S., K.P. Vadrevu, M.S. Mon, and C. Justice. 2021. Contemporary forest loss in Myanmar: effect of democratic transition and subsequent timber bans on landscape structure and composition. *Ambio* 50 (4): 914–928.
- Chen, H., M. Guan, and H. Li. 2021. Air quality prediction based on integrated dual LSTM model. *IEEE Access* 9: 93285–93297.
- Crutzen, P.J., and M.O. Andreae. 1990. Biomass burning in the tropics: impact on atmospheric chemistry and biogeochemical cycles. *Science* 250 (4988): 1669–1678.
- Eaturu, A. and K.P. Vadrevu. (2021). Evaluation of sentinel-3 SLSTR data for mapping fires in forests, peatlands, and croplands—a case study over Australia, Indonesia, and India. In *Biomass burning in South and Southeast Asia* (pp. 39–59). CRC Press.
- FAOSTAT. (2020). <http://www.fao.org/faostat/en/>
- Goldammer, J.G. (ed.). (2012). *Fire in the tropical biota: Ecosystem processes and global challenges* (Vol. 84). Springer Science & Business Media.
- Gupta, P.K., V.K. Prasad, C. Sharma, A.K. Sarkar, Y. Kant, K.V.S. Badarinath, and A.P. Mitra. 2001. CH₄ emissions from biomass burning of shifting cultivation areas of tropical deciduous forests—experimental results from ground-based measurements. *Chemosphere-Global Change Science* 3 (2): 133–143.
- Han, J., G. Kim, C. Lee, Y. Han, U. Hwang, S. Kim. (2019). Predictive models of fire via deep learning exploiting colorific variation. In 2019 international conference on artificial intelligence in information and communication (ICAIIIC), 11–13 Feb. 2019. 579–581.

- Hayasaka, H., I. Noguchi, E.I. Putra, N. Yulianti, and K. Vadrevu. 2014. Peat-fire-related air pollution in Central Kalimantan, Indonesia. *Environmental Pollution* 195: 257–266.
- Hochreiter, S., and J. Schmidhuber. 1997. Long short-term memory. *Neural Computation* 9: 1735–1780.
- Ito, A., and J.E. Penner. 2005. Historical emissions of carbonaceous aerosols from biomass and fossil fuel burning for the period 1870–2000[J]. *Global Biogeochemical Cycles* 19 (2): 1–14.
- Justice, C., G. Gutman, and K.P. Vadrevu. 2015. NASA land cover and land use change (LCLUC): an interdisciplinary research program. *Journal of Environmental Management* 148: 4–9.
- Kadir, E. A., H. Irie, and S.L. Rosa. (2019). Modeling of wireless sensor networks for detection land and forest fire hotspot. In 2019 international conference on electronics, information, and communication (ICEIC), 22–25 Jan 2019 2019. 1–5.
- Kadir, E.A., M. Othman, and S.L. Rosa. (2021). Smart sensor system for detection and forecasting forest fire hotspot in Riau province Indonesia. In 2021 international congress of advanced technology and engineering (ICOTEN), 4–5 July 2021 2021. 1–6.
- Kadir, E.A., S.L. Rosa, and R. A. Ramadhan. (2020). Detection of forest fire used multi sensors system for peatland area in Riau Province. In: W. Sutopo (ed.). 6th international conference on industrial mechanical electrical chemical engineering 2020, 2020 Surakarta, Indonesia. AIP Conference Proceedings.
- Kant, Y., V.K. Prasad, and K.V.S. Badarinath. 2000. Algorithm for detection of active fire zones using NOAA AVHRR data. *Infrared Physics and Technology* 41 (1): 29–34.
- Khabarov, N., E. Moltchanova, and M. Obersteiner. 2008. Valuing weather observation systems for forest fire management. *IEEE Systems Journal* 2: 349–357.
- Kharol, S.K., K.V.S. Badarinath, A.R. Sharma, D.V. Mahalakshmi, D. Singh, and V.K. Prasad. 2012. Black carbon aerosol variations over Patiala city, Punjab, India—a study during agriculture crop residue burning period using ground measurements and satellite data. *Journal of Atmospheric and Solar-Terrestrial Physics* 84: 45–51.
- Kukuk, S.H.B., and Z.H. Kiliminci. (2021). Comprehensive analysis of forest fire detection using deep learning models and conventional machine learning algorithms. *International Journal of Computational and Experimental Science and Engineering (IJCESEN)*. Turkey: Iskender Akkurt.
- Lamjiak, T., R. Kaewthongrach, B. Sirinaovakul, P. Hanpattanakit, A. Chithaisong, and J. Polvichai. 2021. Characterizing and forecasting the responses of tropical forest leaf phenology to El Nino by machine learning algorithms. *PLoS ONE* 16: e0255962.
- Lasko, K., and K. Vadrevu. 2018. Improved rice residue burning emissions estimates: accounting for practice-specific emission factors in air pollution assessments of Vietnam. *Environmental Pollution* 236: 795–806.
- Lasko, K., K.P. Vadrevu, and T.T.N. Nguyen. 2018. Analysis of air pollution over Hanoi, Vietnam using multi-satellite and MERRA reanalysis datasets. *PLoS ONE* 13 (5): e0196629.
- Lasko, K., K.P. Vadrevu, V. Bandaru, T.N.T. Nguyen, and H.Q. Bui. (2021). PM_{2.5} emissions from biomass burning in South/Southeast Asia—Uncertainties and trade-offs. In *Biomass burning in south and Southeast Asia* (pp. 149–169). CRC Press.
- Lasko, K., K.P. Vadrevu, V.T. Tran, E. Ellicott, T.T. Nguyen, H.Q. Bui, and C. Justice. 2017. Satellites may underestimate rice residue and associated burning emissions in Vietnam. *Environmental Research Letters* 12 (8): 085006.
- Le, X.-H., H.V. Ho, G. Lee, and S. Jung. 2019. Application of long short-term memory (LSTM) neural network for flood forecasting. *Water* 11: 1387.
- Liang, H., M. Zhang, and H. Wang. 2019. A neural network model for wildfire scale prediction using meteorological factors. *IEEE Access* 7: 176746–176755.
- Martins, J.A., and M.S. Dias. 2009. The impact of smoke from forest fires on the spectral dispersion of cloud droplet size distributions in the Amazonian region. *Environmental Research Letters* 4 (1): 015002.

- Mohan, K.V.M., A.R. Satish, K.M. Rao, R.K. Yarava, and G.C. Babu. (2021). Leveraging machine learning to predict wild fires. In 2021 2nd international conference on smart electronics and communication (ICOSEC), 7–9 Oct. 2021. 1393–1400.
- NASA 2021. Fires Hotspot Information. 2010–2021. USA.
- Omar, N., A. Al-Zebari, and A. Sengur. (2021). Deep learning approach to predict forest fires using meteorological measurements. In 2021 2nd international informatics and software engineering conference (IISEC), 16–17 Dec. 2021. 1–4.
- Petropoulos, G.P., K.P. Vadrevu, and C. Kalaitzidis. 2013. Spectral angle mapper and object-based classification combined with hyperspectral remote sensing imagery for obtaining land use/cover mapping in a Mediterranean region. *Geocarto International* 28 (2): 114–129.
- Prapas, I., S. Kondylatos, I. Papoutsis, G. Camps-Valls, M. Ronco, M.-N.F. Ndez-Torres, M.P. Guillem, and N. Carvalhais. (2021). Deep learning methods for daily wildfire danger forecasting. In 35th conference on neural information processing systems (NeurIPS 2021). Sydney, Australia.
- Prasad, K.V., and K.V.S. Badarinath. 2006. Soil surface nitrogen losses from agriculture in India: a regional inventory within agroecological zones (2000–2001). *The International Journal of Sustainable Development and World Ecology* 13 (3): 173–182.
- Prasad, V.K., and K.V.S. Badarinth. 2004. Land use changes and trends in human appropriation of above ground net primary production (HANPP) in India (1961–98). *Geographical Journal* 170 (1): 51–63.
- Prasad, V.K., E. Anuradha, and K.V.S. Badarinath. 2005. Climatic controls of vegetation vigor in four contrasting forest types of India—evaluation from National Oceanic and atmospheric administration’s advanced very high resolution radiometer datasets (1990–2000). *International Journal of Biometeorology* 50 (1): 6–16.
- Prasad, V.K., K.V.S. Badarinath, S. Yonemura, and H. Tsuruta. 2004. Regional inventory of soil surface nitrogen balances in Indian agriculture (2000–2001). *Journal of Environmental Management* 73 (3): 209–218.
- Prasad, V.K., Y. Kant, and K.V.S. Badarinath. 2001a. CENTURY ecosystem model application for quantifying vegetation dynamics in shifting cultivation areas: a case study from Rampa forests, Eastern Ghats (India). *Ecological Research* 16 (3): 497–507.
- Prasad, V.K., Y. Kant, and K.V.S. Badarinath. 2002a. Land use changes and modeling carbon fluxes from satellite data. *Advances in Space Research* 30 (11): 2511–2516.
- Prasad, V.K., Y. Kant, P.K. Gupta, C. Elvidge, and K.V.S. Badarinath. 2002b. Biomass burning and related trace gas emissions from tropical dry deciduous forests of India: a study using DMSP-OLS data and ground-based measurements. *International Journal of Remote Sensing* 23 (14): 2837–2851.
- Prasad, V.K., Y. Kant, P.K. Gupta, C. Sharma, A.A. Mitra, and K.V.S. Badarinath. 2001b. Biomass and combustion characteristics of secondary mixed deciduous forests in Eastern Ghats of India. *Atmospheric Environment* 35 (18): 3085–3095.
- Prasad, V.K., M. Lata, and K.V.S. Badarinath. 2003. Trace gas emissions from biomass burning from northeast region in India—estimates from satellite remote sensing data and GIS. *The Environmentalist* 23 (3): 229–236.
- Sigsgaard, T., B. Forsberg, I. Annesi-Maesano, A. Blomberg, A. Bølling, C. Boman, J. Bønløkke, M. Brauer, N. Bruce, M.E. Héroux, and M.R. Hirvonen. 2015. Health impacts of anthropogenic biomass burning in the developed world. *European Respiratory Journal*. 46 (6): 1577–1588.
- Thapa, S.K., J.F. de Jong, A.R. Hof, N. Subedi, L.R. Joshi, and H.H. Prins. 2022. Fire and forage quality: postfire regrowth quality and pyric herbivory in subtropical grasslands of Nepal. *Ecology and Evolution* 12 (4): e8794.
- Szopa, S., D.A. Hauglustaine, and P. Ciais. (2007). Relative contributions of biomass burning emissions and atmospheric transport to carbon monoxide interannual variability. *Geophysical Research Letters*. 34(18).
- Vadrevu, K., and K. Lasko. 2015. Fire regimes and potential bioenergy loss from agricultural lands in the Indo-Gangetic Plains. *Journal of Environmental Management* 148: 10–20.

- Vadrevu, K. 2015. International regional science meeting of NASA-LCLUC: land cover/land use change (LC/LUC) and environmental impacts in South Asia, Coimbatore, India, 19–23 January 2013. *Journal of Environmental Management* 148: 1–163.
- Vadrevu, K., A. Eaturu, E. Casadaban, K. Lasko, W. Schroeder, S. Biswas, L. Giglio, and C. Justice. 2022a. Spatial variations in vegetation fires and emissions in South and Southeast Asia during COVID-19 and pre-pandemic. *Scientific Reports* 12 (1): 1–21.
- Vadrevu, K., A. Heinimann, G. Gutman, and C. Justice. 2019. Remote sensing of land use/cover changes in South and Southeast Asian Countries. *International Journal of Digital Earth* 12 (10): 1099–1102.
- Vadrevu, K., T. Ohara, and C. Justice. 2017. Land cover, land use changes and air pollution in Asia: a synthesis. *Environmental Research Letters* 12 (12): 120201.
- Vadrevu, K.P., and K.V.S. Badarinath. 2009. Spatial pattern analysis of fire events in Central India—a case study. *Geocarto International* 24 (2): 115–131.
- Vadrevu, K.P., and C.O. Justice. 2011. Vegetation fires in the Asian region: satellite observational needs and priorities. *Global Environmental Research* 15 (1): 65–76.
- Vadrevu, K.P. 2008. Analysis of fire events and controlling factors in eastern India using spatial scan and multivariate statistics. *Geografiska Annaler: Series a, Physical Geography* 90 (4): 315–328.
- Vadrevu, K.P., K.V.S. Badarinath, and E. Anuradha. 2008. Spatial patterns in vegetation fires in the Indian region. *Environmental Monitoring and Assessment* 147 (1): 1–13.
- Vadrevu, K.P., A. Eaturu, S. Biswas, K. Lasko, S. Sahu, J.K. Garg, and C. Justice. 2020. Spatial and temporal variations of air pollution over 41 cities of India during the COVID-19 lockdown period. *Scientific Reports* 10 (1): 1–15.
- Vadrevu, K.P., L. Giglio, and C. Justice. 2013. Satellite based analysis of fire–carbon monoxide relationships from forest and agricultural residue burning (2003–2011). *Atmospheric Environment* 64: 179–191.
- Vadrevu, K.P., K. Lasko, L. Giglio, and C. Justice. 2014a. Analysis of Southeast Asian pollution episode during June 2013 using satellite remote sensing datasets. *Environmental Pollution* 195: 245–256.
- Vadrevu, K.P., T. Ohara, and C. Justice, (eds.). (2018). Land-atmospheric research applications in South and Southeast Asia.
- Vadrevu, K.P., T. Ohara, and C. Justice (eds.). (2021a). Biomass burning in south and Southeast Asia: impacts on the biosphere, Volume Two. CRC Press.
- Vadrevu, K.P., T. Ohara, and C. Justice (eds.). (2021b). Biomass burning in South and Southeast Asia: mapping and monitoring, volume one. CRC Press.
- Vadrevu, K.P., T. Ohara, and C. Justice (eds.). (2014a). Air pollution in Asia. *Environmental Pollution* (Barking, Essex: 1987), 195, pp.233–235.
- Vadrevu, T. Le Toan, S.S. Ray, and C. Justice. (2022a). Remote sensing of agriculture and land cover/land use changes in South and Southeast Asian Countries. Springer Cham.
- Vadrevu, K.P., A. Eaturu, E. Casadaban, and S. Biswas. (2022b). Agricultural fires in South Asian countries and implications. In *Remote sensing of agriculture and land cover/land use changes in South and Southeast Asian Countries* (pp. 501–516). Springer, Cham.
- Wooster, M.J., G.J. Roberts, L. Giglio, D.P. Roy, P.H. Freeborn, L. Boschetti, C. Justice, C. Ichoku, W. Schroeder, D. Davies, and A.M. Smith. 2021. Satellite remote sensing of active fires: History and current status, applications and future requirements. *Remote Sensing of Environment* 267: 112694.